


HIV Surveillance by Race/Ethnicity

National Center for HIV/AIDS, Viral Hepatitis, STD & TB Prevention
Division of HIV/AIDS Prevention




Diagnoses of HIV Infection among Adults and Adolescents, by Race/Ethnicity, 2010–2014—United States and 6 Dependent Areas



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.

Rates of Diagnoses of HIV Infection among Adults and Adolescents, by Race/Ethnicity 2010–2014—United States



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.



HIV Infection in Blacks/African Americans United States and 6 Dependent Areas

Total 222,185 diagnoses of HIV infection in 2010–2014

- Blacks/African Americans: 45% of total
- Black/African American women: 62% of women
- Blacks/African Americans: 64% of infections attributed to heterosexual contact^a
- Black/African American children: 64% of children aged <13 years

In 2014, 44% of diagnoses of HIV infection among adults and adolescents were in blacks/African Americans.

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.



HIV Infection in Hispanics/Latinos United States and 6 Dependent Areas

Total 222,185 diagnoses of HIV infection from 2010–2014

- Hispanics/Latinos: 23% of total
- Hispanic/Latino women: 17% of women
- Hispanics/Latinos : 18% of infections attributed to heterosexual contact^a
- Hispanic/Latino children: 11% of children aged <13 years

In 2014, 24% of diagnoses of HIV infection among adults and adolescents were in Hispanics/Latinos.

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting


^a Hispanics/Latinos can be of any race.

^b Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.




Diagnoses of HIV Infection and Population by Race/Ethnicity, 2014—United States

Diagnoses of HIV infection
N = 44,073



Population, United States
N = 318,857,056



American Indian/Alaska Native
Asian
Black/African American

Hispanic/Latino^a
Native Hawaiian/other Pacific Islander
White


Multiple races

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.


^a Hispanics/Latinos can be of any race.

Diagnoses of HIV Infection and Population among Adult and Adolescent Males, by Race/Ethnicity 2014—United States

Diagnoses of HIV Infection
N = 35,571



Male Population, United States
N = 130,043,526



■ American Indian/Alaska Native
■ Asian
■ Black/African American

■ Hispanic/Latino^a
■ Native Hawaiian/Other Pacific Islander
■ White

■ Multiple races


Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.




Diagnoses of HIV Infection and Population among Adult and Adolescent Females, by Race/Ethnicity 2014—United States

Diagnoses of HIV Infection
N = 8,328



Female Population, United States
N = 136,147,401



■ American Indian/Alaska Native
■ Asian
■ Black/African American

■ Hispanic/Latino^a
■ Native Hawaiian/other Pacific Islander
■ White


■ Multiple races

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.


Diagnoses of HIV Infection among Adult and Adolescent Blacks/African Americans, by Sex and Transmission Category 2014—United States and 6 Dependent Areas

Males
N = 14,319



- Male-to-male sexual contact
- Injection drug use (IDU)
- Male-to-male sexual contact and IDU

Females
N = 5,131



- Heterosexual contact^a
- Other^b

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.


^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.




Diagnoses of HIV Infection among Adult and Adolescent Hispanics/Latinos^a, by Sex and Transmission Category 2014—United States and 6 Dependent Areas

Males
N = 9,379



- Male-to-male sexual contact
- Injection drug use (IDU)
- Male-to-male sexual contact and IDU

Females
N = 1,490



- Heterosexual contact^b
- Other^c



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.

^b Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^c Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

Diagnoses of HIV Infection among Adult and Adolescent Whites, by Sex and Transmission Category 2014—United States and 6 Dependent Areas




Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.


^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

Diagnoses of HIV Infection among Adult and Adolescent Hispanics/Latinos^a, by Sex and Place of Birth 2014—United States and 6 Dependent Areas

Males
N = 9,379



Females
N = 1,490



United States
Central America
South America
Cuba
Mexico
Puerto Rico
Other
Unknown

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.



Diagnoses of HIV Infection among Adult and Adolescent Males, by Race/Ethnicity, 2014—United States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	170	18.3
Asian	898	13.4
Black/African American	14,305	94.0
Hispanic/Latino ^a	8,832	41.5
Native Hawaiian/other Pacific Islander	49	22.0
White	10,513	12.6
Multiple races	803	40.6
Total	35,571	27.4

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^a Hispanics/Latinos can be of any race.

^b Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.



Diagnoses of HIV Infection among Adult and Adolescent Females, by Race/Ethnicity, 2014—United States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	49	5.1
Asian	137	1.8
Black/African American	5,128	30.0
Hispanic/Latino ^a	1,350	6.5
Native Hawaiian/Other Pacific Islander	8	3.5
White	1,483	1.7
Multiple races	174	8.2
Total	8,328	6.1


Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^a Hispanics/Latinos can be of any race.

^b Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.



Diagnoses of HIV Infection among Adults and Adolescents, by Region and Race/Ethnicity 2014—United States



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.


^a Hispanics/Latinos can be of any race.



Rates of Diagnoses of HIV Infection among Adult and Adolescent Blacks/African Americans, 2014—United States

N = 19,433

Total Rate = 60.1




Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



Rates of Diagnoses of HIV Infection among Adult and Adolescent Hispanics/Latinos^a, 2014—United States

N = 10,182

Total Rate = 24.2



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.


^a Hispanics/Latinos can be of any race.



Rates of Diagnoses of HIV Infection among Adult and Adolescent Whites, 2014—United States

N = 11,996

Total Rate = 7.0



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



Deaths of Persons with Diagnosed HIV Infection by Race/Ethnicity, 2013—United States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	66	2.8
Asian ^a	64	0.4
Black/African American	7,581	19.4
Hispanic/Latino ^b	2,664	4.9
Native Hawaiian/other Pacific Islander	5	0.9
White	5,028	2.5
Multiple races	874	14.1
Total^c	16,281	5.1

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Deaths of persons with a diagnosed HIV infection may be due to any cause. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.

^c Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.



Adults and Adolescents Living with Diagnosed HIV Infection by Race/Ethnicity, Year-end 2013—United States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	2,862	152.4
Asian ^a	11,462	83.2
Black/African American	397,067	1,243.8
Hispanic/Latino ^b	189,946	462.0
Native Hawaiian/other Pacific Islander	866	201.3
White	297,003	174.0
Multiple races	32,319	815.1
Total^c	931,526	353.2

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.


^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.

^c Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum up to the total column.




Stage 3 (AIDS) Classifications among Racial/Ethnic Minorities with Diagnosed HIV Infection, 1985–2014 United States and 6 Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Includes American Indians/Alaska Natives, Asians, blacks/African Americans, Hispanics/Latinos, and Native Hawaiians/other Pacific Islanders. Hispanics/Latinos can be of any race.



Percentages of Stage 3 (AIDS) Classifications among Adults and Adolescents with Diagnosed HIV Infection, by Race/Ethnicity and Year of Diagnosis, 1985–2014 – United States and 6 Dependent Areas




Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.

^b Includes Asian/Pacific Islander legacy cases.

Rates of Stage 3 (AIDS) Classifications among Adults and Adolescents with Diagnosed HIV Infection, by Race/Ethnicity 2000–2014—United States





Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^a Hispanics/Latinos can be of any race.

^b Includes Asian/Pacific Islander legacy cases.



Stage 3 (AIDS) Classifications and Population, by Race/Ethnicity, 2014—United States



■ American Indian/Alaska Native
■ Asian^b
■ Black/African American

■ Hispanic/Latino^a
■ Native Hawaiian/other Pacific Islander
■ White

■ Multiple races

Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.


^a Hispanics/Latinos can be of any race.

^b Includes Asian/Pacific Islander legacy cases.

Rates of Stage 3 (AIDS) Classifications among Adult and Adolescent Blacks/African Americans with Diagnosed HIV Infection, 2014—United States

N = 9,983

Total Rate = 30.9




Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are not provided for the U.S. dependent areas because U.S. census information for race/ethnicity is limited for these areas.

Rates of Stage 3 (AIDS) Classifications among Adult and Adolescent Hispanics/Latinos^a with Diagnosed HIV Infection, 2014—United States

N = 4,267

Total Rate = 10.1




Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are not provided for the U.S. dependent areas because U.S. census information for race/ethnicity is limited for these areas.

^a Hispanics/Latinos can be of any race.

Rates of Stage 3 (AIDS) Classifications among Adult and Adolescent Whites with Diagnosed HIV Infection, 2014—United States


N = 5,297

Total Rate = 3.1



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are not provided for the U.S. dependent areas because U.S. census information for race/ethnicity is limited for these areas.

Survival after Classification of Stage 3 (AIDS) during 1998–2010, by Months Survived and Race/Ethnicity United States and 6 Dependent Areas



Note. Data exclude persons whose month of diagnosis or month of death is unknown.

^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.



Adults and Adolescents Living with Diagnosed HIV Infection Ever Classified as Stage 3 (AIDS), by Race/Ethnicity, Year-end 2013—United States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	1,506	80.2
Asian ^a	5,997	43.5
Black/African American	217,074	680.0
Hispanic/Latino ^b	111,441	271.1
Native Hawaiian/Other Pacific Islander	468	108.8
White	159,913	93.7
Multiple races	19,730	497.6
Total^c	516,129	195.7

Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population. Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum up to the total column.

^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.

^c Includes persons of unknown race/ethnicity.

